

The global nilpotent variety is Lagrangian

Victor Ginzburg

The purpose of this note is to present a short elementary proof of a theorem due to Faltings and Laumon, saying that the global nilpotent cone is a Lagrangian substack in the cotangent bundle of the moduli space of G -bundles on a complex compact curve. This result plays a crucial role in the Geometric Langlands program, see [BD], since it insures that the $\mathcal{D}$ -modules on the moduli space of G -bundles whose characteristic variety is contained in the global nilpotent cone are automatically *holonomic*, hence, e.g. have finite length.

Let (M, ω) be a smooth symplectic algebraic variety. A (possibly singular) algebraic subvariety $Y \subset M$ is said to be isotropic, resp. Lagrangian, if the tangent space, $T_y Y$, at any regular point $y \in Y$ is an isotropic, resp. Lagrangian, vector subspace in the symplectic vector space $T_y M$ (we always assume Y to be *reduced*, but not necessarily irreducible). The following characterisation of isotropic subvarieties proved, e.g. in [CG, Prop. 1.3.30], will be used later : *$Y \subset M$ is isotropic if and only if for any smooth locally closed subvariety $W \subset Y$, we have $\omega|_W = 0$* (here W is possibly contained in the singular locus of Y). An advantage of this characterisation is that it allows to extend the notion of ‘being isotropic’ from algebraic subvarieties to semi-algebraic constructible subsets. Thus, we call a semi-algebraic constructible subset $Y \subset M$ isotropic if $\omega|_W = 0$ for any smooth locally closed algebraic variety $W \subset Y$.

Now, let M be a smooth stack that can be locally presented as $p : \widetilde{M} \rightarrow M$, where $\widetilde{M}$ is a smooth algebraic variety and p is a smooth surjective morphism; e.g., M is locally isomorphic to the quotient of a smooth algebraic variety modulo an algebraic action of an algebraic group, see e.g. [LMB]. We have a natural diagram

$$T^*M \leftarrow T^*M \times_M \widetilde{M} \xrightarrow{\epsilon} T^*\widetilde{M}.$$

A substack $Z \subset T^*M$ is said to be constructible, resp. isotropic or Lagrangian, if $\epsilon(Z \times_M \widetilde{M})$ is a constructible, resp. isotropic or Lagrangian,

subset of $T^*\widetilde{M}$ relative to the standard symplectic structure on the cotangent bundle of a smooth variety.

Let X be a smooth complex compact connected algebraic curve of genus $g > 1$, and G a complex semisimple¹ group. Below, we write Bun_G for the moduli space of principal algebraic G -bundles on X , regarded as a *stack*, cf. [LMB], rather than a scheme. In particular, no stability conditions on principal bundles $P \in Bun_G$ are imposed. Given a principal G -bundle P , let $\mathfrak{g}_P$ and $\mathfrak{g}_P^*$ denote the associated vector bundles corresponding to the adjoint and co-adjoint representations of G , respectively. Let $\mathbf{P}$ be the universal bundle on $Bun_G \times X$, and $p : Bun_G \times X \rightarrow Bun_G$ the projection.

The cotangent stack, T^*Bun_G , is a stack, see [BD, n.1.1.1], which is relatively representable over Bun_G by the affine spectrum of the sheaf of algebras $\mathrm{Sym}(R^1p_*\mathfrak{g}_P)$, the Symmetric algebra of the first derived pushforward sheaf. Note that for all $i > 1$, we have: $R^ip_*\mathfrak{g}_P = 0$, since $\dim X = 1$. Hence the formation $R^1p_*(-)$ is right-exact, and therefore commutes with base change. For a scheme S and a morphism: $S \rightarrow Bun_G$, write $\mathbf{P}(S)$ for the pull-back of the universal bundle to $S \times X$. Using the base change, one obtains the following (stack version of the) Kodaira-Spencer formula for the set of S -points of the fiber of T^*Bun_G over $\mathbf{P}(S)$:

$$\begin{aligned} T_{\mathbf{P}(S)}^*Bun_G &= \Gamma(S, \mathrm{Hom}_{\mathrm{Coh}}(R^1p_*\mathfrak{g}_{\mathbf{P}(S)}, \mathcal{O}_S)) \\ &= \Gamma(S, \mathrm{Hom}_{D^b(\mathrm{Coh})}(Rp_*\mathfrak{g}_{\mathbf{P}(S)}[1], \mathcal{O}_S)) \\ &= \Gamma(S, \mathrm{Hom}_{D^b(\mathrm{Coh})}(\mathfrak{g}_{\mathbf{P}(S)}, Rp^!\mathcal{O}_S[-1])) \\ &= \Gamma(S \times X, \mathfrak{g}_{\mathbf{P}(S)}^* \otimes \Omega_{X \times S/S}^1) \simeq \Gamma(S \times X, \mathfrak{g}_{\mathbf{P}(S)} \otimes \Omega_{X \times S/S}^1), \end{aligned} \tag{1}$$

where the second isomorphism exploits the fact that the complex $Rp_*\mathfrak{g}_{\mathbf{P}(S)}[1]$ is concentrated in non-positive degrees, and the last isomorphism uses the identification $\mathfrak{g}_P^* \simeq \mathfrak{g}_P$ induced by the Killing form on $\mathfrak{g}$.

Write $\mathcal{N}$ for the nilpotent cone in $\mathfrak{g}$, the zero variety of the set of $\mathrm{Ad} G$ -invariant polynomials on $\mathfrak{g}$ without constant term. Choose a Borel subgroup $B \subset G$ with Lie algebra $\mathfrak{b}$, and let $\mathfrak{n}$ denote the nilradical of $\mathfrak{b}$. Kostant proved [Ko], see also [CG, ch.6], that $\mathcal{N}$ is equal, as a subscheme of $\mathfrak{g}$, to the image of the Springer resolution, the morphism: $G \times_B \mathfrak{n} \rightarrow \mathfrak{g}$, given by the assignment: $(g, x) \mapsto \mathrm{Ad} g(x)$.

¹it is not hard to extend our results to any reductive group, but that would lead to unpleasant dimension shifts in various formulas below, so we restrict ourselves to the semisimple case.

Given a scheme S and a G -bundle P on $S \times X$, we choose local trivialisations of the vector bundles $\mathfrak{g}_P$ and Ω_X^1 and view a local section x of $\mathfrak{g}_P \otimes \Omega_X^1$ as a function $S \times X \rightarrow \mathfrak{g}$. The section x is called *nilpotent* if the corresponding function gives a morphism $S \times X \rightarrow \mathcal{N} \subset \mathfrak{g}$. The notion of a nilpotent section does not depend on the choices of trivialisations involved. Following Laumon [La], define the *global nilpotent cone* as a closed (non-reduced) substack $\mathcal{N}ilp \subset T^*Bun_G$ whose set of S -points is

$$\mathcal{N}ilp(S) = \{(P, x) \mid x \in \Gamma(S \times X, \mathfrak{g}_P \otimes \Omega_X^1), \text{ } x \text{ is nilpotent section}\},$$

where P runs over G -bundles on $S \times X$.

Main Theorem. *$\mathcal{N}ilp$ is a Lagrangian substack in T^*Bun_G .*

Remark. This theorem was first proved, in the special case $G = SL_n$, by Laumon [La]. His argument cannot be generalized to arbitrary semisimple groups. In the general case, the theorem was proved by Faltings [Fa, theorem II.5]. The proof below seems to be more elementary than that of Faltings; it is based on nothing but a few general results of Symplectic geometry. Another proof of the theorem is given in [BD]. That proof is more complicated; however, it potentially leads to a description of the irreducible components of $\mathcal{N}ilp$.

We begin with a few general lemmas. Let (M_1, ω_1) and (M_2, ω_2) be complex algebraic symplectic manifolds, and $\text{pr}_i : M_1 \times M_2 \rightarrow M_i$, the projections. We regard $M_1 \times M_2$ as a symplectic manifold with symplectic form $\text{pr}_1^*\omega_1 - \text{pr}_2^*\omega_2$, involving the minus-sign on the second factor. The following result is a special case of [CG, Prop. 2.7.51].

Lemma 2. *Let $\Lambda_1 \subset M_1$ and $\Lambda \subset M_1 \times M_2$ be smooth algebraic isotropic subvarieties. Then $\text{pr}_2(\text{pr}_1^{-1}(\Lambda_1) \cap \Lambda) \subset M_2$ is an isotropic subvariety.*

Proof. Set $Y := \text{pr}_1^{-1}(\Lambda_1) \cap \Lambda$. Simple linear algebra shows that, for any $y \in Y$, the image of the tangent map $(\text{pr}_2)_* : T_y Y \rightarrow T_{\text{pr}_2(y)} M_2$ is isotropic. We use the characterization of isotropic subvarieties mentioned at the beginning of the paper. Let $W \subset \Lambda_2 := \text{pr}_2(Y)$ be an irreducible smooth subvariety. Observe that the map $\text{pr}_2 : \text{pr}_2^{-1}(W) \cap Y \rightarrow W$ is *surjective*. Hence, there exists a non-empty smooth Zariski-open dense subset $Y' \subset (\text{pr}_2^{-1}(W) \cap Y)_{\text{red}}$ such that the restriction $\text{pr}_2 : Y' \rightarrow W$ has surjective differential at any point of Y' . Therefore the tangent space at the generic point of W is isotropic. Whence the tangent space at every point of

W is isotropic, by continuity. It follows that any smooth subvariety of Λ_2 is isotropic, and the lemma follows. $\square$

Given a manifold N we write λ_N for the canonical 1-form on T^*N , usually denoted ‘ pdq ’, such that $d\lambda$ is the canonical symplectic 2-form on T^*N . Let $f : N_1 \rightarrow N_2$ be a morphism of smooth algebraic varieties. Identify $T^*(N_1 \times N_2)$ with $T^*N_1 \times T^*N_2$ via the standard map multiplied by (-1) on the factor T^*N_2 . The canonical 1-form on $T^*(N_1 \times N_2)$ becomes under the above identification equal to $\text{pr}_1^*(\lambda_{N_1}) - \text{pr}_2^*(\lambda_{N_2})$. We endow $T^*N_1 \times T^*N_2$ with the corresponding symplectic form $\text{pr}_1^*(d\lambda_{N_1}) - \text{pr}_2^*(d\lambda_{N_2})$; it is induced from the canonical symplectic form on $T^*(N_1 \times N_2)$ via the identification.

Introduce the following closed subvariety

$$Y_f = \{(n_1, \alpha_1), (n_2, \alpha_2) \in T^*N_1 \times T^*N_2 \mid n_2 = f(n_1), \alpha_1 = 0 = f^*(\alpha_2)\}. \quad (3)$$

Lemma 4. *The image of Y_f under the second projection $\text{pr}_2 : T^*N_1 \times T^*N_2 \rightarrow T^*N_2$ is an isotropic subvariety in T^*N_2 .*

Proof. Using the above explained identification of $T^*(N_1 \times N_2)$ with $T^*N_1 \times T^*N_2$ involving a sign, the conormal bundle to the graph of f can be written as the subvariety

$$\Lambda = \{(n_1, \alpha_1), (n_2, \alpha_2) \in T^*N_1 \times T^*N_2 \mid n_2 = f(n_1), \alpha_1 = f^*(\alpha_2)\}.$$

Observe that the canonical 1-form $\text{pr}_1^*(\lambda_{N_1}) - \text{pr}_2^*(\lambda_{N_2})$ on $T^*N_1 \times T^*N_2$ vanishes identically on Λ . Hence, Λ is an isotropic subvariety, and we may apply Lemma 1 to $M_1 = T^*N_1$, $M_2 = T^*N_2$, and $\Lambda_1 = T_{N_1}^*N_1 = \text{zero-section}$, and to Λ above. Observe now that we have by definition, $Y_f = \Lambda \cap \text{pr}_1^{-1}(T_{N_1}^*N_1)$. Hence by Lemma 2 the subvariety $\text{pr}_2(Y_f)$ is isotropic. $\square$

Lemma 5. *If N_1 and N_2 are smooth algebraic stacks, and $f : N_1 \rightarrow N_2$ is a representable morphism of finite type, then the assertion of Lemma 4 remains valid.*

Proof. Due to locality of the claim we may (and will) assume N_2 is quasi-compact. Let $\tilde{N}_2$ be a smooth algebraic variety and $\tilde{N}_2 \rightarrow N_2$ a smooth surjective equidimensional morphism. Set $\tilde{N}_1 := \tilde{N}_2 \times_{N_2} N_1$. Note that

the set $Y_f \subset T^*N_1 \times T^*N_2$ defined in (3) may be viewed as a subset in $T^*N_2 \times_{N_2} N_1$. Therefore we have

$$Y_f \times_{N_2} \tilde{N}_2 \subset T^*N_2 \times_{N_2} \tilde{N}_2 \times_{N_2} N_1 = T^*\tilde{N}_2 \times_{N_2} \tilde{N}_1.$$

We must show that the image of $Y_f \times_{N_2} \tilde{N}_2$ is an isotropic subvariety in $T^*\tilde{N}_2$. Let $F : \tilde{N}_1 \rightarrow \tilde{N}_2$ be the natural morphism, and $Y_F \subset T^*\tilde{N}_2 \times_{\tilde{N}_2} \tilde{N}_1$ the corresponding subvariety of Lemma 4. Observe that $T^*\tilde{N}_2 \times_{\tilde{N}_2} \tilde{N}_1 = T^*\tilde{N}_2 \times_{N_2} N_1$ and $Y_f \times_{N_2} \tilde{N}_2 = Y_F$. Hence, Lemma 4 applied to F shows that the image of $Y_F \times_{N_2} \tilde{N}_2$ in $T^*\tilde{N}_2$ is isotropic. The claim follows. $\square$

Choose a Borel subgroup $B \subset G$ with Lie algebra $\mathfrak{b}$, and let $\mathfrak{n}$ denote the nilradical of $\mathfrak{b}$. Given a field K of characteristic zero, write $G(K), B(K), \mathfrak{b}(K)$, etc., for the corresponding sets of K -rational points. The following result seems to be well-known; it is included here for the reader's convenience.

Lemma 6. *For any field $K \supset \mathbb{C}$ and any $x \in \mathcal{N}(K)$, there exists an element $g \in G(K)$ such that $\text{Ad } g(x) \in \mathfrak{n}(K)$.*

Proof. Since the Jacobson-Morozov theorem holds for any field of characteristic zero, one may find an $\mathfrak{sl}_2$ -triple $(x, h, x^-) \subset \mathfrak{g}(K)$ associated to the given nilpotent element $x \in \mathfrak{g}(K)$. The eigenspaces of the semisimple endomorphism $\text{ad } h : \mathfrak{g} \rightarrow \mathfrak{g}$ corresponding to non-negative eigenvalues span a parabolic subalgebra $\mathfrak{p}_x \subset \mathfrak{g}$, which is defined over K . Writing $\mathfrak{u}_x$ for the nilradical of $\mathfrak{p}_x$, by construction, we have: $x \in \mathfrak{u}_x(K)$. Clearly, if $\mathfrak{b}_x \subset \mathfrak{p}_x$, is a Borel subalgebra defined over K , and $\mathfrak{n}_x$ is its nilradical, then $x \in \mathfrak{u}_x(K) \subset \mathfrak{n}_x(K)$.

Thus, it suffices to prove that the parabolic $\mathfrak{p}_x$ contains a Borel subalgebra defined over K . To this end, let $\mathcal{P}$ denote the partial flag variety of all parabolics in $\mathfrak{g}$ of type $\mathfrak{p}_x$. There is a unique $\mathfrak{p} \in \mathcal{P}(K)$ such that $\mathfrak{p} \supset \mathfrak{b}$, where $\mathfrak{b}$ is our fixed Borel subalgebra. Now, the group $G(K)$ acts transitively on $\mathcal{P}(K)$ (this follows easily from the Bruhat decomposition, see [Ja]). We deduce that there exists $g \in G(K)$ such that $\text{Ad } g(\mathfrak{p}) = \mathfrak{p}_x$. But then, $\text{Ad } g(\mathfrak{b}) \subset \mathfrak{p}_x$ is a Borel subalgebra defined over K , and we are done. $\square$

Let $f_1, \dots, f_r$, ($r = \text{rk } \mathfrak{g}$), be a set of homogeneous free generators of $\mathbb{C}[\mathfrak{g}]^G$, the algebra of G -invariant polynomials on $\mathfrak{g}$. Let $d_i = \deg f_i$ be the *exponents* of $\mathfrak{g}$. Following Hitchin [Hi] we put:

$$\text{Hitch} := \Gamma(X, \Omega_X^{\otimes d_1}) \bigoplus \dots \bigoplus \Gamma(X, \Omega_X^{\otimes d_r}).$$

This is an affine space of dimension equal to $\dim Bun_G$ (at this point it is used, see [Hi], that genus X is > 1). Hitchin has defined a morphism $\pi : T^*Bun_G \rightarrow Hitch$, by assigning to any pair $(s, P) \in T^*Bun_G$, where $P \in Bun_G$ and $s \in T_P^*Bun_G \simeq \Gamma(X, \mathfrak{g}_{P(s)} \otimes \Omega_X^1)$, see (1), the element $\pi(s, P) = \oplus_{i=1}^r f_i(s) \in Hitch$. It is immediate from the construction that the global nilpotent variety is the fiber of π over the zero element $0 \in Hitch$.

Remark. Hitchin actually worked in the setup of *stable* Higgs bundles and not in the setup of stacks. But his construction of the map π extends to the stack setup verbatim. We make no use of any additional properties of the map π established in [Hi].

Lemma 7. *Nilp is an isotropic substack in T^*Bun_G .*

Proof. Write $N_2 = Bun_B$ for the moduli stack of principal B -bundles. By an old result of Harder [Ha], any G -bundle on a curve has a B -reduction, hence the natural morphism of stacks $f : Bun_B \rightarrow Bun_G$ is surjective.

Let P be an algebraic G -bundle on the curve X , and s a nilpotent regular section of $\mathfrak{g}_P \otimes \Omega_X^1$. Harder showed further, using a key rationality result of Steinberg [St], that the bundle P is locally-trivial in the Zariski topology. Thus, trivializing P on the generic point of X , one may identify the restriction of s to the generic point with a nilpotent element of $\mathfrak{g}(K)$, where $K = \mathbb{C}(X)$ is the field of rational functions on X . Hence Lemma 6 implies that there exists a B -reduction of P over the generic point of X such that $s \in \mathfrak{n}_P \otimes \Omega_X^1$. Here, a B -reduction is a section of the associated bundle $B \backslash P$. The fibers of $B \backslash P$ being projective varieties (isomorphic to $B \backslash G$), any section of $B \backslash P$ defined over the generic point of X extends to the whole of X . Thus, there exists a B -reduction over X of the G -bundle P such that $s \in \mathfrak{n}_P \otimes \Omega_X^1$.

Further, let $k \supset \mathbb{C}$ be a field, and $S = \text{Spec}(k)$. For any $P \in Bun_B(k)$, we have:

$$T_P^*Bun_B = H^1(X, \mathfrak{b}_P)^* = H^0(X, \mathfrak{b}_P^* \otimes \Omega_X^1) = H^0(X, (\mathfrak{g}_P/\mathfrak{n}_P) \otimes \Omega_X^1).$$

It follows that in the notation of Lemma 5, for $N_1 = Bun_B$ and $N_2 = Bun_G$, we have $\mathcal{N}ilp = \text{pr}_2(Y_f)$. Observe further that Bun_B is the union of a countable family of open substacks of finite type over Bun_G each. Thus, Lemma 5 implies that $\mathcal{N}ilp$ is the union of a *countable* family of isotropic substacks.

We claim that if an algebraic stack S is the union of a countable family $\{S_i\}_{i \in \mathbb{N}}$ of locally closed substacks, then any quasi-compact substack of S can be covered by finitely many S_i 's. To prove this, we may assume without loss of generality that S is itself quasi-compact. Choose a smooth surjective morphism $\tilde{S} \twoheadrightarrow S$, where $\tilde{S}$ is a scheme of finite type. By our assumptions, there exists a countable family $\{\tilde{S}_i\}_{i \in \mathbb{N}}$ of locally closed subschemes of $\tilde{S}$ such that: $\tilde{S} = \cup_i \tilde{S}_i$. Hence, there exists $n \gg 0$ such that $\tilde{S}$ equals the union of the closures of $\tilde{S}_1, \dots, \tilde{S}_n$, thanks to Baer theorem. Hence, $\tilde{S}_1 \cup \dots \cup \tilde{S}_n$ is Zariski dense in $\tilde{S}$, and $\dim(\tilde{S} \setminus (\tilde{S}_1 \cup \dots \cup \tilde{S}_n)) < \dim \tilde{S}$. Arguing by induction on $\dim \tilde{S}$ we deduce that $\tilde{S}$ is covered by finitely many $\tilde{S}_i$'s. Hence, for any field $k \supset \mathbb{C}$, the quasi-compact set $S(k)$ is covered by finitely many subsets $S_i(k)$.

Thus, we have proved that any open quasi-compact substack of $\mathcal{N}ilp$ can be covered by finitely many isotropic substacks. This implies that $\mathcal{N}ilp$ is itself isotropic. $\square$

Proposition 8. $\dim \mathcal{N}ilp = \dim Bun_G$.

Proof. We observe that, since Bun_G is an equi-dimensional smooth stack, each irreducible component of T^*Bun_G has dimension $\geq 2 \dim Bun_G$. To see this, one may replace Bun_G by an open quasi-compact substack Y , which admits a smooth surjective morphism $p : \tilde{Y} \twoheadrightarrow Y$, where $\tilde{Y}$ is a smooth algebraic variety and fibers of p are purely m -dimensional. Then, $T^*Y \times_Y \tilde{Y}$ is a closed subscheme of $T^*\tilde{Y}$ locally defined by m equations. Hence, for each irreducible component T_j^* of T^*Y we find:

$$\dim T_j^* = \dim(T_j^* \times_Y \tilde{Y}) - m \geq (\dim T^*\tilde{Y} - m) - m = 2(\dim \tilde{Y} - m) = 2 \dim Y.$$

It follows that each irreducible component of any fiber of the Hitchin morphism $\pi : T^*Bun_G \rightarrow Hitch$ has dimension $\geq 2 \dim Bun_G - \dim Hitch = \dim Bun_G$. But we have proved that each component of $\mathcal{N}ilp = \pi^{-1}(0)$ is an isotropic subvariety. Thus, $\dim \mathcal{N}ilp = \dim Bun_G$. $\square$

Remark. Although the inequality: $\dim T^*Bun_G \geq 2 \dim Bun_G$ is no longer true if X has genus one or zero, it has been shown in [BD] that the stack $\mathcal{N}ilp$ still has pure dimension equal to $\dim Bun_G$.

Corollary 9. *Every irreducible component of any fiber of π has dimension equal to $\dim Bun_G$. In particular, the Hitchin morphism π is flat.*

Proof. There is a natural $\mathbb{C}^*$ -action on $Hitch$, such that $t \in \mathbb{C}^*$ acts on the direct summand $\Gamma(X, \Omega_X^{\otimes d_i}) \subset Hitch$ via multiplication by t^{d_i} . The map $\pi : T^*Bun_G \rightarrow Hitch$ is $\mathbb{C}^*$ -equivariant relative to the above-defined $\mathbb{C}^*$ -action on $Hitch$ and the standard $\mathbb{C}^*$ -action on T^*Bun_G by dilations along the fibers, respectively. Clearly, zero is the only fixed point of the $\mathbb{C}^*$ -action on $Hitch$ and, moreover, it is contained in the closure of any other $\mathbb{C}^*$ -orbit on $Hitch$. It follows, since dimension of any irreducible component of any fiber $\leq$ dimension of the special fiber, that for any $h \in Hitch$ we have: $\dim \pi^{-1}(h) \leq \dim \pi^{-1}(0)$. But for curves of genus > 1 , Proposition 8 yields: $\dim \pi^{-1}(0) = \dim Nilp = \dim Bun_G$. Thus, for any $h \in Hitch$ we get: $\dim \pi^{-1}(h) \leq \dim Bun_G$. On the other hand, the dimension of each irreducible component of any fiber of the morphism $\pi : T^*Bun_G \rightarrow Hitch$ is no less than $\dim T^*Bun_G - \dim Hitch = \dim Bun_G$. This proves the opposite inequality. $\square$

The proof of the main theorem is completed by the following stronger result

Theorem 10. *The stack T^*Bun_G is a local complete intersection, and $Nilp$ is a Lagrangian complete intersection in T^*Bun_G .*

Proof. The claims being local, we may replace Bun_G by an open quasi-compact substack Y which admits a presentation of the form $p : \tilde{Y} \twoheadrightarrow Y$, where $\tilde{Y}$ is a smooth algebraic variety and p is a smooth surjective morphism with fibers of pure dimension m . As we have observed in the proof of Proposition 8, $T^*Y \times_Y \tilde{Y}$ is a closed subscheme of $T^*\tilde{Y}$ locally defined by m equations and, moreover, $\dim T^*Y \geq 2 \dim Bun_G$. On the other hand, since fibers of the Hitchin map π have dimension $= \dim Bun_G$, we find: $\dim T^*Y \leq \dim Bun_G + \dim Hitch = 2 \dim Bun_G$. This proves that the stack T^*Bun_G is a local complete intersection.

Further, $Nilp$ being the zero fiber of the surjective morphism π , it is defined by $\dim Bun_G$ equations in T^*Bun_G . The equality $\dim Nilp = \dim Bun_G$ (Proposition 8) implies that $Nilp$ is a complete intersection in T^*Bun_G . Finally, since $Nilp$ is equidimensional (Corollary 9), Lemma 7 implies that $Nilp$ is Lagrangian. $\square$

I am grateful to V. Drinfeld for his invaluable help, and also for his extreme patience while answering my numerous foolish questions.

References

- [BD] Beilinson A., Drinfeld V.: *Quantization of Hitchin's Integrable system and Hecke eigen-sheaves*. Manuscript available at the following address
<http://www.math.uchicago.edu/~benzvi>
- [CG] Chriss N., Ginzburg V.: *Representation Theory and Complex Geometry*, Birkhäuser Boston, 1997.
- [Fa] Faltings G.: *Stable G-bundles and projective connections*, J. Algebraic Geom. **2** (1993), 507–568.
- [Ha] Harder G.: *Halbeinfache Gruppenschemata über Dedekindringen*, Invent. Math., **4** (1967), 165–191 .
- [Hi] Hitchin N.: *Stable bundles and integrable systems*, Duke Math. J., **54** (1987), 91–114.
- [Ja] Jantzen J.C.: *Representations of algebraic groups*. Pure and Applied Mathematics, **131**. Academic Press, 1987.
- [Ko] Kostant B.: *Lie group representations on polynomial ring*, Amer. J. Math., **85** (1963), 327–404.
- [LMB] Laumon G., Moret-Bailly L.: *Champs algébriques*, *Ergebnisse der Mathematik* **39**, Springer, Berlin-NY, 1999.
- [La] Laumon G.: *Une analogue global du cône nilpotent*, Duke Math. J., **57** (1988), 647–671.
- [St] Steinberg R.: *Regular elements of semisimple algebraic groups*, Publ. Math. IHES, **25** (1965), 281–312.

V.G.: University of Chicago, Department of Mathematics, Chicago, IL 60637, USA;
ginzburg@math.uchicago.edu